

1-Low Dose—Handy Pain Relief

Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg	Pain reliever

Uses for the temporary relief of minor aches and pains or as recommended by your doctor.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ asthma (wheezing) ■ facial swelling ■ shock

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers.

Aspirin may cause stomach bleeding.

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if you have

■ asthma ■ ulcers ■ stomach problems (such as heartburn, upset stomach or stomach pain) that persist or recur ■ bleeding problems

Ask a doctor or pharmacist before use if you are

taking a prescription drug for anticoagulation (blood thinning), gout, diabetes, or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- new symptoms occur
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. IT IS
ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3
MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY
A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR
COMPLICATIONS DURING DELIVERY.

Keep out of the reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours, not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at room temperature 15°-30° C (50°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients black iron oxide, colloidal silicon dioxide, corn starch, D&C yellow 10, FD&C yellow #6, hypromellose, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

2-Stop Sneezing

Drug Facts

Active ingredient (in each tablet)

Loratadine 10 mg

Purpose

Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies ■runny nose ■sneezing ■itchy, watery eyes ■itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- adults and children 6 years and over: 1 tablet daily, not more than 1 tablet in 24 hours
- children under 6years: ask a doctor
- consumers with liver or kidney disease: ask a doctor

Other information

- do not use if blister unit is broken or torn**
- store at room temperature 20°-25° C (68°-77°F)
- protect from excessive moisture

Inactive ingredients lactose monohydrate, magnesium stearate, povidone, pregelatinized starch

Questions or comments 1-800-555-5555

3-Sparkling White Toothpaste

Drug Facts

Active ingredient

Sodium fluoride 0.243% Anticavity toothpaste

Purpose

Uses

helps protect against cavities

Warnings

Keep out of reach of children under 6 yrs. of age. If more than used for brushing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 2 yrs. & older: brush teeth thoroughly after meals or at least twice a day or use as directed by a dentist
- do not swallow
- to minimize swallowing use a pea-sized amount in children under 6
- supervise children's brushing until good habits are established
- children under 2 yrs.: ask a dentist

Inactive ingredients

glycerin, hydrated silica, water, sorbitol, sodium hexametaphosphate, propylene glycol, flavor, PEG-12, cocamidopropyl betaine, sodium lauryl sulfate, carbomer, sodium saccharin, poloxamer 407, polyethylene oxide, xanthan gum, sodium hydroxide, cellulose gum, titanium dioxide

Questions? 1-800-555-5555

4-Motion Sickness Relief

Drug Facts	
Active ingredient (in each tablet)	Purpose
Meclizine HCl 25 mg	
Uses for the prevention and treatment of the nausea, vomiting or dizziness associated with motion sickness	
Warnings	
■Do not give to children under 12 years of age unless directed by a doctor	
Ask a doctor before use if you have	
■breathing problems such as emphysema or chronic bronchitis	
■glaucoma	
■difficulty in urination due to enlargement of the prostate gland	
Ask a doctor or pharmacist before use if you are	
taking sedatives or tranquilizers	
When using this product ■drowsiness may occur	
■avoid alcoholic beverages	
■alcohol, sedatives and tranquilizers may increase drowsiness	
■use caution when driving a motor vehicle or operating machinery	
If pregnant or breast-feeding , ask a health professional before use.	
Keep out of reach of children . In case of overdose, get medical help or contact a Poison Control Center right away	
Directions	
To prevent motion sickness, the first dose should be taken one hour before starting activity	
■ Adults and children 12 years and over : take 1 to 2 tablets once daily, or as directed by a doctor. Do not exceed 2 tablets in 24 hours. Not for frequent or prolonged use except on the advice of a doctor.	
■ Do not exceed recommended dosage.	
Other information	
■store below 86°F (30°C) ■protect from heat & humidity	
Inactive ingredients anhydrous lactose, colloidal silicon dioxide, cornstarch, D&C yellow #10, magnesium stearate, microcrystalline cellulose, sodium starch glycolate.	

5-Stomach Relief

Drug Facts	
Active ingredient (per tablet)	Purpose
Calcium carbonate USP750 mg	
Uses relieves ■heartburn ■sour stomach ■acid indigestion ■upset stomach associated with these symptoms	
Warnings	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.	
When using this product ■do not take more than 10 tablets in 24 hours ■do not use the maximum dosage for more than 2 weeks	
Keep out of reach of children	
Directions	
■chew 2-4 tablets as symptoms occur, or as directed by a doctor	
Other information	
■for calcium/magnesium/sodium content see Supplement Facts	
■store below 25° C (77°F). Keep the container tightly closed.	
Inactive ingredients calcium carbonate, sorbitol, dextrose, sucrose, cocoa powder, microcrystalline cellulose, natural and artificial flavors, magnesium stearate, corn starch, maltodextrin, guar gum, silicon dioxide. Contains: milk, soy.	
Safety sealed – do not use if printed inner seal beneath cap is missing or broken	
(Questions? 1-800-555-5555 weekdays)	

6-All-purpose Cold Medicine

Drug Facts

Active ingredient (in each caplet)

Active ingredient (in each caplet)	Purpose
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg.....	Antihistamine
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses ■ temporarily relieves these common cold symptoms:

- minor aches and pains ■ headache ■ sore throat ■ nasal congestion
- cough ■ sneezing and runny nose ■ helps clear nasal passages ■ relieves cough to help you sleep ■ temporarily reduces fever

Warnings

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage.

Sore Throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other product containing acetaminophen ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease

- high blood pressure ■ thyroid disease ■ diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ do not exceed recommended dosage

- excitability may occur, especially in children ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur

- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new

symptoms occur ■ cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-555-5555) Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
adults and children 12 years and over: ■ take 2 caplets every 4 hours
- swallow whole – do not crush, chew or dissolve ■ do not take more than 12 caplets in 24 hours
- children under 12 years: do not use this adult product in children under 12 years of age, this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

- store at room temperature 20°-35° C (68°-77°F)
- do not use if carton is opened or if blister unit is broken
- see side panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, carnauba wax, corn starch, FD&C blue #1 aluminum lake, flavors, hypromellose, iron oxide, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, potassium sorbate, powdered cellulose, pregelatinized starch, propylene glycol shellac, sodium benzoate, sodium citrate, sodium starch glycolate, sucralose, titanium dioxide

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

7-Flu Plus

Drug Facts

Active ingredient (in each tablet)

Aspirin 500 mg Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg Antihistamine
Dextromethorphan hydrobromide 15 mg Cough suppressant

Uses ■ temporarily relieves these symptoms due to a cold or flu:
■ minor aches and pains ■ headache ■ coughing ■ sore throat
■ runny nose ■ sneezing ■ temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers should not use this medicine for chickenpox or flu-like symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ in children under 12 years of age

Ask a doctor before use if you have ■ ulcers ■ glaucoma
■ cough with excessive phlegm (mucus) ■ breathing problems such as emphysema, chronic bronchitis or asthma ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
■ stomach problems that persist or recur ■ difficulty in urination due to enlargement of the prostate gland ■ a sodium-restricted diet

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking a prescription drug for anticoagulation (blood thinning), diabetes, gout, or arthritis

When using this product ■ **do not exceed recommended dosage** ■ excitability may occur, especially in children

■ may cause marked drowsiness ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition ■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

■ adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 6 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
■ children under 12 years: do not use.

Other information

■ each tablet contains: sodium 387 milligrams
■ Phenylketonurics: Contains Phenylalanine 6.7 mg per tablet
■ store at room temperature. Avoid excessive heat.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, calcium silicate, dimethylpolysiloxane, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, saccharin sodium, sodium benzoate, sodium bicarbonate

Questions or comments? 1-800-555-5555 (Mon-Fri AM-5PM EST)

8-Indigestion Help

Drug Facts

Active ingredient (in each tablet)	Purpose
---------------------------------------	---------

Anhydrous citric acid 1000 mg.....	Antacid
Aspirin 325 mg	Analgesic
Sodium bicarbonate (heat-treated) 1700 mg.....	Antacid

Uses for the relief of ■ heartburn acid indigestion, and sour stomach when accompanied with headache or body ache and pains
■ upset stomach with headache from overindulgence in food or drink
■ pain alone (headache or body and muscular aches and pains)

Warnings

Reye's syndrome: Children and teenagers should not use this medicine for chickenpox or flu-like symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding.

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if you have

■ asthma ■ ulcers ■ bleeding problems ■ stomach problems that last or come back frequently, such as heartburn, upset stomach, or pain ■ a sodium restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (blood thinning), diabetes, gout, or arthritis

When using this product do not exceed recommended dosage

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ symptoms get worse or last more than 10 days ■ redness or swelling is present ■ ringing in the ears or loss of hearing occurs ■ new symptoms occur

Drug Facts (continued)

If pregnant or breast-feeding ask a health professional before use. It is especially important not to use aspirin during the last 3 months

of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away

Directions ■ fully dissolve 2 tablets in 4 ounces of water before taking

■ adults and children 12 years and over: 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 8 tablets in 24 hours

■ adults 60 years and over: 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 4 tablets in 24 hours

■ children under 12 years: consult a doctor

Other information

■ each tablet contains: sodium 504 milligrams

■ phenylketonurics: Contains phenylalanine 9 mg per tablet

■ store at room temperature. Avoid excessive heat.

Inactive ingredients aspartame, docusate sodium, flavor, povidone, sodium benzoate

Questions or comments? 1-800-555-5555 (Mon-Fri 9AM-5PM EST)

9-Pain Be Gone

Drug Facts

Active ingredient (in each caplet)	Purpose
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....	Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug
Uses ■temporarily relieves minor aches and pains due to: ■minor pain of arthritis ■muscular aches ■backache ■menstrual cramps ■headache ■toothache ■the common cold ■temporarily reduces fever	
Warnings	
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■hives ■facial swelling ■asthma (wheezing) ■shock ■skin reddening ■rash ■blisters If an allergic reaction occurs, stop use and seek medical help right away.	
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■are age 60 or older ■have had stomach ulcers or bleeding problems ■take a blood thinning (anticoagulant) or steroid drug ■take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■have 3 or more alcoholic drinks every day while using this product ■take more or for a longer time than directed	
Do not use ■if you have ever had an allergic reaction to any other pain reliever/fever reducer ■right before or after heart surgery	
Ask a doctor before use if you have ■problems or serious side effects from taking pain relievers or fever reducers ■stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ulcers ■bleeding problems ■high blood pressure ■heart or kidney disease ■taken a diuretic ■reached age 60 or older	
Ask a doctor or pharmacist before use if you are ■taking any other drug containing an NSAID (prescription or nonprescription) ■taking a blood thinning (anticoagulant) or steroid drug ■under a doctor's care for any serious condition ■taking any other drug	

When using this product ■take with food or milk if stomach upset occurs ■long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if ■you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■pain gets worse or lasts more than 10 days ■fever gets worse or lasts more than 3 days ■stomach pain or upset gets worse or lasts ■redness or swelling is present in the painful area ■any new symptoms appear ■you have difficulty swallowing ■it feels like the capsule is stuck in your throat ■you develop heartburn

If pregnant or breast-feeding, ask a health professional before use. It is especially important to not use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions ■do not take more than directed ■the smallest effective dose should be used ■do not take longer than 10 days, unless directed by a doctor (see Warnings) ■drink a glass full of water with each dose ■if taken with food, this product may take longer to work

adults and children 12 years and older: ■take 1 capsule every 8 to 12 hours while symptoms last ■for the first dose you may take 2 capsules within the first hour ■do not exceed 2 capsules in any 8-12 hour period ■do not exceed 3 capsules in a 24-hour period
children under 12 years: ■ask a doctor

Other information ■each capsule contains: sodium 20mg ■store at 20°-25° C (68°-77°F) avoid high humidity and excessive heat above 40°C (104°F) ■read all directions and warnings before use

Inactive ingredients: FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-800-555-5555 (Mon-Fri 9AM-5PM EST)

10-Crampless Wonder Relief

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Caffeine 60 mg.....	Stimulant
Pyrilamine maleate 15 mg.....	Diuretic

Uses for the temporary relief of these symptoms associated with menstrual periods: ■cramps ■bloating ■water-weight gain ■breast tenderness ■headache ■backache ■muscle aches ■fatigue

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■more than 8 caplets in 24 hours, which is the maximum daily amount ■with other drugs containing acetaminophen ■3 or more alcoholic drinks every day while using this product

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, as a doctor or pharmacist.

Ask a doctor before use if you have ■ liver disease ■glaucoma ■difficulty in urination due to enlargement of the prostate gland ■a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■taking sedatives or tranquilizers

When using this product ■you may get drowsy ■avoid alcoholic drinks ■excitability may occur, especially in children ■alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■limit the use of caffeine-containing medication, foods or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if ■new symptoms occur ■redness or swelling is present ■pain gets worse or lasts more than 10 days

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before

use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults, as well as for children, even if you do not notice any symptoms.

Directions ■do not take more than the recommended dose ■adults and children 12 years and older: take 2 caplets with water. Repeat every 6 hours, as needed. Do not exceed 8 caplets per day. ■children under 12 years: consult a doctor

Other information ■store at room temperature

Inactive ingredients: carnauba wax, croscarmellose sodium, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, propylene glycol, shellac, titanium dioxide, tricetin

Questions or comments? 1-800-555-5555 (Mon-Fri 9AM-5PM EST)

11-Crampless PM

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg.....	Pain reliever
Diphenhydramine citrate 38 mg.....	Nighttime sleep aid

Uses for relief of occasional sleeplessness when associated with minor aches and pains from premenstrual and menstrual periods (dysmenorrhea).

Warnings

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage.

Do not use ■ with any other product containing acetaminophen ■ in children under 12 years of age ■ with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness. ■ pain gets worse or lasts more than 10 days ■ new symptoms occur ■ redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

■ do not take more than the recommended dose (see **overdose warning**)
■ adults and children 12 years and older: take 2 caplets at bedtime if needed, or as directed by a doctor ■ children under 12 years: do not use

Other information ■ store at room temperature

Drug Facts (continued)

Inactive ingredients carnauba wax, corn starch, croscarmellose sodium, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, hypromellose, microcrystalline cellulose, povidone, propylene glycol, shellac, stearic acid, titanium dioxide

Questions or comments? 1-800-555-5555

(Mon – Fri 9AM – 5PM EST)

12-Adult Cough & Cold Syrup

Drug Facts

Active ingredient (in each 5 mL tsp) Purposes

Dextromethorphan HBr, USP 10 mg.....	Cough suppressant
Guaifenesin, USP 100 mg.....	Expectorant
Phenylephrine HCl , USP 5 mg.....	Nasal decongestant

Uses ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes ■ temporarily relieves these symptoms occurring with a cold:
■ nasal congestion ■ cough due to minor throat and bronchial irritation

Warnings

Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ trouble urinating due to an enlarged prostate gland
■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product ■ do not exceed recommended dose

Stop use and ask a doctor if ■you get nervous, dizzy, or sleepless
■symptoms do not get better within 7 days or are accompanied by fever ■cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

■do not take more than 6 doses in any 24-hour period ■this adult product is not intended for use in children under 12 years of age ■**adults and children 12 years and over:** 2 teaspoons every 4 hours; **children under 12 years:** do not use

Other information

■each teaspoon contains: sodium 3 mg
■store between 20° - 25° C (68° - 77°F)
■dosage cup provided

Drug Facts (cont'd)

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 9AM to 5PM EST at
1-800-555-5555

13-Nighty Night Cold/Flu

Drug Facts

Active ingredient (in each 15 mL TBSP)

Acetaminophen 500 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 15 mg.....Cough suppressant
Doxylamine succinate 6.25 mgAntihistamine

Purpose

Uses

■ temporarily relieves common cold/flu symptoms:
■ headache ■ sore throat ■ fever ■ runny nose ■ cough due to minor throat and bronchial irritation ■ sneezing ■ minor aches and pains

Warnings

Overdose Warning: Taking more than the recommended dose can cause serious health problems, including liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers /fever reducers. Acetaminophen may cause liver damage.

Do not use ■ if you are now taking a prescription monamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ with any other product containing acetaminophen

Ask a doctor before use if you have ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ cough that lasts or is chronic such as occurs with smoking, asthma or emphysema ■ a sodium restricted diet

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ avoid alcoholic drinks ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ new symptoms appear ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be

signs of a serious condition. ■ redness or swelling is present ■ sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions

■ do not use more than directed (see overdose warning)
■ adults (12 years and older): 2 TBSP (30 mL) in dose cup provided ■ repeat every 6 hours; not more than 4 doses per day ■ children under 12: ask a doctor

Other information ■ each TBSP contains: sodium 18 mg
■ store at 68°- 77°F (20°- 25°C)

Inactive ingredients alcohol 10%, citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

14-Headache Relief PM

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 500mg.....	Pain reliever
Diphenhydramine HCl 25 mg..... (equiv. to Diphenhydramine citrate 38 mg)	Nighttime sleep-aid

Purpose

Uses ■ for the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness:

Warnings

Overdose warning: Taking more than the recommended dose can cause serious health problems, including liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/ fever reducers. Acetaminophen may cause liver damage.

Do not use ■ in children under 12 years of age ■ with any other product containing acetaminophen
■ with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery. This product will cause drowsiness.

Stop use and ask a doctor if ■ redness or swelling is present ■ new symptoms occur ■ pain gets worse or lasts for more than 10 days ■ sleeplessness lasts continuously for more than 2 weeks.

Insomnia may be a symptom of serious underlying medical illness. ■ fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

■ do not use more than directed (see overdose warning) ■ adults and children 12 years and over: take 2 tablets at bedtime, if needed. ■ do not use this adult product in children under 12 years of age.

Inactive ingredients carnauba wax, crospovidone, FD&C blue #1, FD&C blue #2, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Store at room temperature (59°- 86°F)

15-Severe Cold

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg.....Antihistamine/cough suppressant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

■ temporarily relieves these symptoms of the common cold:

■ runny nose ■ sneezing ■ headache ■ minor aches and pains ■ nasal congestion ■ cough ■ sore throat ■ temporarily reduces fever

Warnings

Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage.

Sore Throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with another product containing acetaminophen ■ with any other product containing diphenhydramine, even one used on skin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure
■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland
■ cough accompanied by too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma or emphysema ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product ■ do not exceed recommended dose

■ excitability may occur, especially in children ■ marked drowsiness may occur
■ alcohol, sedatives and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur
■ pain, cough or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-555-5555) Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over: ■ take 2 caplets every 4 hours ■ do not take more than 12 caplets in 24 hours
- children 6 to under 12 years of age: take 1 caplet every 4 hours ■ do not take more than 5 caplets in 24 hours
- children under 6 years: do not use this adult product in children under 6 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

- store between 20° - 25° C (68° - 77°F)
- do not use if carton is opened or if blister unit is broken
- see side panel for lot number and expiration date

Inactive ingredients carnauba wax, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium citrate, sodium starch glycolate, talc, titanium dioxide

Questions or comments? call 1-888-555-5555

16-Ache Prevent

Drug Facts

Active ingredient (per tablet) Purposes

Aspirin (NSAID)*400 mg.....Pain reliever
Caffeine 32 mgPain reliever aid

*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:

- headache ■ a cold ■ muscular aches ■ arthritis ■ toothache ■ menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have ■ asthma ■ ulcers ■ bleeding problems ■ stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ high blood pressure ■ heart or kidney disease ■ taken a diuretic ■ reached age 60 or older

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ new symptoms occur ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding ■ stomach pain or upset gets worse or lasts ■ ringing in the ears or loss of hearing occurs ■ painful area is red

or swollen ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not use more than directed (see overdose warning) ■ drink a full glass of water with each dose ■ adults and children over 12 years of age and over: take 2 tablets every 6 hours; not more than 8 tablets in 24 hours, or as directed by a doctor. ■ children under 12: consult a doctor

Other information

Keep carton for full drug facts. Read all warnings and directions before use. Store at 20° – 25° C (68° – 77° F). Protect from moisture.

Inactive ingredients hypromellose, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate, starch

Questions? Call 1-800-555-5555 or visit our website at www.acheprevent.com

17-Aspirin

Drug Facts

Active ingredient (in each tablet) Purposes

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- provides temporary relief of
■ headache ■ pain and fever of colds ■ muscle pain ■ menstrual pain
■ toothache ■ minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if you have ■ stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ bleeding problems ■ ulcers ■ asthma ■ high blood pressure ■ heart or kidney disease ■ taken a diuretic ■ reached age 60 or over

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur ■ ringing in the ears or loss of hearing occurs

- stomach pain or upset gets worse or lasts

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of a pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
■ adults: 1-2 tablets with water. Dosage may be repeated every 4 hours, not to exceed 12 tablets in 24 hours ■ **children under 12:** consult a doctor

Other information ■ store at room temperature

Inactive ingredients hypromellose, starch, titanium dioxide

Questions? Call toll free **1-800-555-5555**

18-Ibuprofen

Drug Facts

Active ingredient (in each brown tablet) Purpose

Ibuprofen USP, 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
- headache ■ muscular aches ■ toothache ■ minor pain of arthritis ■ menstrual cramps ■ backache ■ the common cold ■ temporarily reduces fever

Warnings

Allergy Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) ■ rash ■ blisters ■ skin reddening. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ are age 60 or older

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if you have ■ bleeding problems ■ high blood pressure ■ ulcers ■ stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ problems or serious side effects from taking pain relievers or fever reducers ■ heart or kidney disease ■ taken a diuretic ■ reached age 60 or older

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs ■ long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if ■ redness or swelling is present in the painful area ■ you feel faint, vomit blood or have bloody or black stools. These are signs of stomach bleeding ■ fever gets worse or lasts more than 3 days ■ stomach pain or upset gets worse or lasts ■ any new symptoms appear ■ pain gets worse or lasts

more than 10 days

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed ■ the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at controlled room temperature
- avoid excessive heat 40° C (104° F) ■ use by expiration date on package

Inactive ingredients carnauba wax, cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions? Call 1-800-STORE or www.store.com

19 Extra Strength Acetaminophen

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500mg.....	Pain reliever/fever reducer
Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ arthritis ■ cold ■ toothache ■ flu ■ premenstrual and menstrual cramps ■ reduces fever	
Warnings	
Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/ fever reducers. Acetaminophen may cause liver damage.	
Do not use ■ with any other product containing acetaminophen	
Stop use and ask a doctor if ■ new symptoms occur ■ redness or swelling is present ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
Directions	
■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets every 4-6 hours as needed. Do not take more than 8 caplets in 24 hours ■ Children under 12 years: do not use this adult Extra Strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) and could cause serious health problems	
Other information ■ do not use if inner foil seal printed "sealed for your protection" is broken or missing ■ store at room temperature between 15° and 30°C (59° and 86°F) ■ keep tightly closed to protect from moisture	
Inactive ingredients hypromellose. May also contain: carnauba wax, cellulose, corn starch, hydroxypropyl cellulose, magnesium stearate, polyethylene glycol, polysorbate 80, polyvinyl alcohol, polyvinylpyrrolidone, povidone, silicon dioxide, sodium carboxymethylcellulose, sodium starch glycolate, starch, stearic acid, talc, titanium dioxide, triacetin	
Questions? Call 1-800-774-2678	

20-Pain Relieving Patch

Drug Facts

Active ingredient (in each tablet)	Purpose
Camphor 1.2%.....	Topical Analgesic
Menthol 5.7%.....	Topical Analgesic
Methyl Salicylate 6.3%.....	Topical Analgesic

Uses ■ For temporary relief of minor aches & pains of muscles & joints associated with: ■ arthritis ■ simple backache ■ sprains ■ bruises ■ strains

Warnings For external use only.

Do not use ■ on wounds or damaged skin. ■ if you are allergic to aspirin or salicylates. ■ with a heating pad. ■ with, or at the same times as, other external analgesic products. ■ if you are allergic to any ingredients of this product.

When using this product ■ do not use otherwise than as directed. ■ avoid contact with the eyes, mucous membranes or rashes

Stop use and ask a doctor if ■ rash, itching or excessive skin irritation develops. ■ conditions worsen. ■ symptoms persist for more than 7 days. ■ symptoms clear up and occur again within a few days.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions **Adults and children 12 years of age and over:** ■ Clean and dry affected area. ■ Remove patch from film. ■ Apply to affected area not more than 3 to 4 times daily for 7 days. ■ Remove patch from the skin after at most 8 hours' application. **Children under 12 years of age:** ■ Consult a doctor.

Other information

- Avoid storing product in direct sunlight.
- Protect product from excessive moisture.

Inactive ingredients Butylated Hydroxytoluene, Hydrogenated Rosin Glycerol Ester, Maleated Rosin Glycerin Ester, Natural Rubber, Perfume, Polybutene, Polyisobutylene, Silicon Dioxide, Starch Grafted Acrylate, Titanium Dioxide, Tocopherol Acetate, Zinc Oxide