

2-Stop Sneezing

Drug Facts	
Active ingredient (in each tablet) Medicine 10 mg	Purpose Antihistamine
Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies ■runny nose ■sneezing ■itchy, watery eyes ■itching of the nose or throat	
Warnings Do not use if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose. When using this product do not take more than directed. Taking more than directed may cause drowsiness. Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away. If pregnant or breast-feeding , ask a health professional before use. Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away	
Directions ■adults and children 6 years and over: 1 tablet daily, not more than 1 tablet in 24 hours ■children under 6years: ask a doctor ■consumers with liver or kidney disease: ask a doctor	
Other information ■do not use if blister unit is broken or torn ■store at room temperature 20°-25° C (68°-77° F) ■protect from excessive moisture	
Inactive ingredients lactose monohydrate, magnesium stearate, povidone, pregelatinized starch	
Questions or comments 1-800-555-5555	

3-All-purpose Cold Relief

Drug Facts	
Active ingredient (in each caplet)	Purpose
Medicine 325 mg.....	Pain reliever/fever reducer
Medicine 2 mg.....	Antihistamine
Medicine 10 mg.....	Cough suppressant
Medicine 5 mg.....	Nasal decongestant
Uses ■ temporarily relieves these common cold symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ cough ■ sneezing and runny nose ■ helps clear nasal passages ■ relieves cough to help you sleep ■ temporarily reduces fever	
Warnings Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers or fever reducers. Acetaminophen may cause liver damage. Sore Throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use ■ with any other product containing acetaminophen ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma or emphysema ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery	
Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new	

symptoms occur ■ cough comes back or occurs with rash or headache that lasts
 These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-555-5555) Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years and over: ■ take 2 caplets every 4 hours

■ swallow whole – do not crush, chew or dissolve ■ do not take more than 12 caplets in 24 hours

■ children under 12 years: do not use this adult product in children under 12 years of age, this will provide more than the recommended dose (overdose) and may cause liver damage

Other information

■ store at room temperature 20°-35° C (68°-77° F)

■ do not use if carton is opened or if blister unit is broken

■ see side panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, carnauba wax, corn starch, FD&C blue #1 aluminum lake, flavors, hypromellose, iron oxide, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, potassium sorbate, powdered cellulose, pregelatinized starch, propylene glycol shellac, sodium benzoate, sodium citrate, sodium starch glycolate, sucralose, titanium dioxide

Questions or comments? call 1-877-895-3665 (English) or 1-888-466-8746 (Spanish)

4-Motion Sickness Relief

Drug Facts	
Active ingredient (in each tablet) Medicine I 25 mg	Purpose .Antiemetic
Uses for the prevention and treatment of the nausea, vomiting or dizziness associated with motion sickness	
Warnings <ul style="list-style-type: none"> ■Do not give to children under 12 years of age unless directed by a doctor Ask a doctor before use if you have <ul style="list-style-type: none"> ■breathing problems such as emphysema or chronic bronchitis ■glaucoma ■difficulty in urination due to enlargement of the prostate gland 	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product ■drowsiness may occur <ul style="list-style-type: none"> ■avoid alcoholic beverages ■alcohol, sedatives and tranquilizers may increase drowsiness ■use caution when driving a motor vehicle or operating machinery 	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away	
Directions To prevent motion sickness, the first dose should be taken one hour before starting activity <ul style="list-style-type: none"> ■Adults and children 12 years and over: take 1 to 2 tablets once daily, or as directed by a doctor. Do not exceed 2 tablets in 24 hours. Not for frequent or prolonged use except on the advice of a doctor. ■Do not exceed recommended dosage. 	
Other information <ul style="list-style-type: none"> ■store below 86°F (30°C) ■protect from heat & humidity 	
Inactive ingredients anhydrous lactose, colloidal silicon dioxide, cornstarch, D&C yellow #10, magnesium stearate, microcrystalline cellulose, sodium starch glycolate.	

5-Stomach Relief

Drug Facts	
Active ingredient (per tablet) Medicine USP750 mg	Purpose ...Antacid
Uses relieves ■heartburn ■sour stomach ■acid indigestion <ul style="list-style-type: none"> ■upset stomach associated with these symptoms 	
Warnings Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.	
When using this product ■do not take more than 10 tablets in 24 hours ■do not use the maximum dosage for more than 2 weeks	
Keep out of reach of children	
Directions <ul style="list-style-type: none"> ■chew 2-4 tablets as symptoms occur, or as directed by a doctor 	
Other information <ul style="list-style-type: none"> ■for calcium/magnesium/sodium content see Supplement Facts ■store below 25° C (77°F). Keep the container tightly closed. 	
Inactive ingredients calcium carbonate, sorbitol, dextrose, sucrose, cocoa powder, microcrystalline cellulose, natural and artificial flavors, magnesium stearate, corn starch, maltodextrin, guar gum, silicon dioxide. Contains: milk, soy.	
Safety sealed – do not use if printed inner seal beneath cap is missing or broken	
Questions? 1-800-555-5555 weekdays	

6-Flu Plus

Drug Facts	
Active ingredient (in each tablet)	Purpose
Medicine 500 mg	..Pain reliever/fever reducer
Medicine 2 mg	Antihistamine
Medicine 15 mg	Cough suppressant
Uses ■temporarily relieves these symptoms due to a cold or flu: ■minor aches and pains ■headache ■coughing ■sore throat ■runny nose ■sneezing ■temporarily reduces fever	
Warnings Reye's syndrome: Children and teenagers should not use this medicine for chickenpox or flu-like symptoms before a doctor is consulted about Reye's syndrome, a rare but serious illness reported to be associated with aspirin. Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■hives ■facial swelling ■asthma (wheezing) ■shock Alcohol Warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers/fever reducers. Aspirin may cause stomach bleeding. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use to sedate children.	
Do not use ■if you are allergic to aspirin or any other pain reliever/fever reducer ■if you are now taking a prescription monamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■in children under 12 years of age	
Ask a doctor before use if you have ■ulcers ■glaucoma ■cough with excessive phlegm (mucus) ■breathing problems such as emphysema, chronic bronchitis or asthma ■persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■stomach problems that persist or recur ■difficulty in urination due to enlargement of the prostate gland ■a sodium-restricted diet	

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are ■taking sedatives or tranquilizers ■taking a prescription drug for anticoagulation (blood thinning), diabetes, gout, or arthritis

When using this product ■do not exceed recommended dosage ■excitability may occur, especially in children
■may cause marked drowsiness ■avoid alcoholic drinks ■alcohol, sedatives, and tranquilizers may increase drowsiness ■be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■an allergic reaction occurs. Seek medical help right away. ■pain or cough gets worse or lasts more than 7 days ■fever gets worse or lasts more than 3 days
■redness or swelling is present ■new symptoms occur ■cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition ■ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

■adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 6 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
■children under 12 years: do not use.

Other information

■each tablet contains: sodium 387 milligrams
■Phenylketonurics: Contains Phenylalanine 6.7 mg per tablet
■store at room temperature. Avoid excessive heat.

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, calcium silicate, dimethylpolysiloxane, docusate sodium, FD&C red #40, FD&C yellow #6, flavors, mannitol, povidone, saccharin sodium, sodium benzoate, sodium bicarbonate

Questions or comments? 1-800-555-5555 (Mon-Fri AM-5PM EST)